



# Nevada

## Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,\* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


| Age group    |             |             |
|--------------|-------------|-------------|
| 15-17        | 18-19       | 15-19       |
| 15th highest | 4th highest | 9th highest |

\*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




# Sexually transmitted diseases Nevada

## Gonorrhea rates, 1982–1992



## Gonorrhea rates, 1992



## National gonorrhea rate ranking,\* 1992


Males – 19th highest    Females – 26th highest

\*National ranking refers to males and females aged 10–24 years.




## Chlamydia cases and rates,\* 1988–1992


Cases




Rates




Cases



Rates



## Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



\*Chlamydia cases and rates refer to males and females of all ages. Data were unavailable for 1990.

†Missing bar indicates fewer than six cumulative cases.



# Related risk behaviors,\* 1993

Nevada

Had sexual intercourse (lifetime)



Had sexual intercourse with four or more people (lifetime)



Had sexual intercourse during the past three months (currently sexually active)




Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



\*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 Nevada Youth Risk Behavior Survey (risk behavior data).

